

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x) g(y)$  USING  
SEPARATION OF VARIABLES**

**1** Use the method of separation of variables to find the general solution of each of the differential equations below. Where reasonable, express the family of solutions as explicit functions of  $x$ .

**(a)**  $(x^2 + 4) \frac{dy}{dx} = 2xy$

**(b)**  $\frac{dy}{dx} = \frac{2y}{x}$

**(c)**  $\frac{dy}{dx} = (1 + y^2) \sqrt{x}$

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x) g(y)$  USING  
SEPARATION OF VARIABLES**

**1** Use the method of separation of variables to find the general solution of each of the differential equations below. Where reasonable, express the family of solutions as explicit functions of  $x$ .

(e)  $(1 + x^2) \frac{dy}{dx} = xy$       (f)  $e^y \cos x - \frac{dy}{dx} \sin^2 x = 0$       (g)  $(\sec x) y' + y^2 = 0$

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x) g(y)$  USING  
SEPARATION OF VARIABLES**

**2** Find the particular solution of  $e^{-x^2} yy' + xy = 0$ ,  $y(0) = 1$ .

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x) g(y)$  USING  
SEPARATION OF VARIABLES**

**3** Find the equation of each graph:

- (a) The graph passes through  $(1, 2)$  and has a slope  $\frac{3y}{x^2}$  at each point  $(x, y)$ .
- (b) The gradient of the tangent at point  $(x, y)$  on a graph is given by  $\frac{-2y}{x}$  and the graph passes through the point  $(1, 2)$ .

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x)g(y)$  USING  
SEPARATION OF VARIABLES**

**5** Consider the differential equation  $\frac{dy}{dx} = 3x^2 \cos^2 y$ .

- (a) Find the particular solution  $y = f(x)$  to the differential equation, satisfying the initial condition  $f(0) = \frac{\pi}{4}$ .
- (b) State the domain and range of the solution found in part (a).

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x) g(y)$  USING  
SEPARATION OF VARIABLES**

- 6** An insect population  $P$  experiences a seasonal growth rate given by  $\frac{dP}{dt} = \frac{\pi}{12} \sin\left(\frac{\pi}{6}t\right)P$ ,  $P(0) = 1$ , where  $P$  is measured in millions and  $t$  is the number of months since the beginning of spring.

Express the time variation of the insect population  $P$  and sketch this variation over the course of one year.

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x) g(y)$  USING  
SEPARATION OF VARIABLES**

**10** Consider the differential equation  $\frac{dy}{dx} = -\frac{2xy}{\log_e y}$ ,  $y > 0$ .

- (a) Find the general solution  $g(x, y) = c$  of this differential equation as an implicit relation between  $x$  and  $y$ , using the substitution  $u = \log_e y$  to complete the integration.
- (b) Find the particular solution passing through the point  $(0, e)$ .
- (c) Explain why  $x = 1$  cannot exist in the solution to part (b).

**SOLVING DIFFERENTIAL EQUATIONS OF THE FORM  $dy/dx = f(x)g(y)$  USING  
SEPARATION OF VARIABLES**

**14** A space probe is launched vertically upwards from the surface of a spherical planet with a radius  $R$ . If the atmospheric drag is ignored, the upwards velocity ( $v \text{ m s}^{-1}$ ) of the probe at height  $h$  metres above the surface of the planet is modelled by the solution of the differential equation  $\frac{dv}{dh} = -\frac{gR^2}{v(R+h)^2}$ ,  $v = u$ , where  $h = 0$  and  $g$  is the gravitational acceleration on the surface of the planet.

(a) Show that  $v^2 = u^2 - \frac{2gR}{1 + \frac{R}{h}}$ .

(b) Hence find the minimum launch velocity  $u$  for the probe to escape the planet's gravity.